

# \***Prevantics**®

Prevent the spread of HAIs with the first and **ONLY** 3.15% CHG/70% IPA formulation with up to 7 days of continued antimicrobial activity.\*\*



## THE CHALLENGE

There were an estimated 722,000 HAIs in U.S. hospitals in 2011, resulting in an estimated 75,000 patient deaths.<sup>1</sup> HAIs affect patients, their families and the people who care for them. They also have an impact on hospital costs and quality. The CDC estimated that CLABSIs cost facilities \$45,816 per infection.<sup>2</sup>

## THE SOLUTION

**Prevantics** products, designed with 3.15% Chlorhexidine Gluconate (w/v) and 70% Isopropyl Alcohol (v/v) formulation, help prevent the spread of infections and help improve patient outcomes.



### **Designed for Convenience**

**Prevantics** products are a convenient delivery system that has no sharp glass to break or activation required.

#### **Prevantics Applicators**

- No glass to break
- Easy to open, easy to use
- Pre-saturated
- Swab: Intuitive prep-pad design
- Swabsticks: Unique 2-sided applicator for interdigital areas and skin folds



Prevantics 1.0 mL Swab



Prevantics 5.1 mL Maxi Swabstick

<sup>1</sup>CDC HAI Prevalence Survey. Magill SS, Edwards JR, Bamberg W, et al. Multistate Point-Prevalence Survey of Health Care–Associated Infections. N Engl J Med 2014;370:1198-208. <sup>2</sup>NCBI Article: Haddadin Y, Regunath H. Central Line Associated Blood Stream Infections (CLABSI). Update March 28, 2017

# **Prevantics** products are the first and only FDA approved 3.15% Chlorhexidine Gluconate (w/v) and 70% Isopropyl Alcohol (v/v) solution.

#### The **Prevantics** Difference


The efficacy of **Prevantics** products closely correlate with dressing change protocols at many facilities.



### **Clinical Case Study**

The following facility has successfully implemented **Prevantics** products to improve patient outcomes.

## The Rush-Copley Medical Center Study resulted in a CLABSI reduction of 59% in year 1 and a 100% reduction by year 4.



At Rush-Copley Medical Center they adjusted practice protocols for infection prevention and Vascular Access with **Prevantics** products to reduce the incidence of CLABSI rates in their facility.<sup>3</sup>

\*\*Swabstick and Maxi Swabstick only

<sup>3</sup>Matocha et al. A Process Improvement Approach to the Elimination of Central Line Associated Bloodstream Infections. Rush-Copley Medical Center, Aurora, IL. October 2011.

#### Prevantics products are available in a variety of convenient, ready-to-use applicators: Swab, Swabstick and Maxi Swabstick.

#### **PRODUCT BENEFITS**

Saves Time 30 SCRUB Records Records	<ul> <li>Pre-saturated. No waiting for activation</li> <li>Unique 2-sided applicators**</li> <li>No glass</li> <li>Easy to open packaging</li> </ul>					
Saves Money	• Effective skin antisepsis will help improve patient outcomes and reduce healthcare costs					
Improves Outcomes	<ul> <li>Up to 7 days of continued antimicrobial activity**</li> <li>Clinical compendium with summary of independent studies</li> <li>8 independent clinical studies demonstrating clinical efficacy with <b>Prevantics</b> products</li> </ul>					
Meets Evidence-Based Clinical Guidelines	Prevantics products meet the skin antisepsis guidelines from:     Image: Stream S					

#### **ORDER INFORMATION**

Individual Boxes	Reorder No.	Swabs/Swabsticks Per Box	Boxes Per Case	Treatment Area Covered
Prevantics Swab 1.0 mL Preinjection Indication	B10800	100	10	2.5" x 2.5" in <sup>2</sup> (6.25 in <sup>2</sup> / 6 x 6 cm)
<b>Prevantics</b> Swabstick 1.6 mL Preinjection and Preoperative Indications	S40750	50	10	Dry Site: 4" x 4" in² (16 in² / 10 x 10 cm) Moist Site: 3" x 5" in² (15 in² / 7.5 x 12.5 cm)
<b>Prevantics</b> Maxi Swabstick 5.1 mL Preinjection and Preoperative Indications	S41950	30	10	Dry Site: 7" x 7" in² (49 in² / 18 x 18 cm) Moist Site: 3" x 7.5" in² (22.5 in² / 7.5 x 19 cm)

Available for Kits	Reorder No.	Swabs/Swabsticks Per Box	Boxes Per Case	Treatment Area Covered
<b>Prevantics</b> Swab 1.0 mL Preinjection Indication	B11400	3000	1	2.5" x 2.5" in <sup>2</sup> (6.25 in <sup>2</sup> / 6 x 6 cm)
<b>Prevantics</b> Swabstick 1.6 mL Preinjection and Preoperative Indications	\$32450	500	1	Dry Site: 4" x 4" in² (16 in² / 10 x 10 cm) Moist Site: 3" x 5" in² (15 in² / 7.5 x 12.5 cm)
<b>Prevantics</b> Compact Swabstick 1.6 mL Preinjection and Preoperative Indications	S42850	500	1	Dry Site: 4" x 4" in² (16 in² / 10 x 10 cm) Moist Site: 3" x 5" in² (15 in² / 7.5 x 12.5 cm)
<b>Prevantics</b> Maxi Swabstick 5.1 mL Preinjection and Preoperative Indications	S27350	300	1	Dry Site: 7" x 7" in <sup>2</sup> (49 in <sup>2</sup> / 18 x 18 cm) Moist Site: 3" x 7.5" in <sup>2</sup> (22.5 in <sup>2</sup> / 7.5 x 19 cm)

\*For dry sites only \*\*Swabstick and Maxi Swabstick only

